



Smoking and Bone Health

**National Institutes of Health
Osteoporosis and Related
Bone Diseases
National Resource Center**

2 AMS Circle
Bethesda, MD 20892-3676

Phone: 202-223-0344
Toll free: 800-624-BONE
TTY: 202-466-4315
Fax: 202-293-2356

Website: www.bones.nih.gov
Email: NIHBoneInfo@mail.nih.gov

The NIH Osteoporosis and Related Bone Diseases National Resource Center is supported by the National Institute of Arthritis and Musculoskeletal and Skin Diseases with contributions from the National Institute on Aging, the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, the National Institute of Dental and Craniofacial Research, the National Institute of Diabetes and Digestive and Kidney Diseases, the NIH Office of Research on Women's Health, and the HHS Office on Women's Health.

The National Institutes of Health (NIH) is a component of the U.S. Department of Health and Human Services (HHS).

May 2016

Many of the health problems caused by tobacco use are well known. Cigarette smoking causes heart disease, lung and esophageal cancer, and chronic lung disease. Additionally, several research studies have identified smoking as a risk factor for osteoporosis and bone fracture.

Facts About Osteoporosis

Osteoporosis is a condition in which bones weaken and are more likely to fracture. Fractures from osteoporosis can result in pain and disability. In the United States, more than 53 million people either already have osteoporosis or are at high risk due to low bone mass. According to the Centers for Disease Control and Prevention, more than 16 million Americans are living with a disease caused by smoking.

In addition to smoking, risk factors for developing osteoporosis include:

- thinness or small frame
- family history of the disease
- being postmenopausal and particularly having had early menopause
- abnormal absence of menstrual periods (amenorrhea)
- prolonged use of certain medications, such as those used to treat lupus, asthma, thyroid deficiencies, and seizures
- low calcium intake
- lack of physical activity
- excessive alcohol intake.

Osteoporosis can often be prevented. It is known as a "silent" disease because, if undetected, bone loss can progress for many years without symptoms until a fracture occurs. It has been called a childhood disease with old age consequences because building healthy bones in youth helps prevent osteoporosis and fractures later in life. However, it is never too late to adopt new habits for healthy bones.

Smoking and Osteoporosis

Cigarette smoking was first identified as a risk factor for osteoporosis decades ago. Studies have shown a direct relationship between tobacco use and

decreased bone density. Analyzing the impact of cigarette smoking on bone health is complicated. It is hard to determine whether a decrease in bone density is due to smoking itself or to other risk factors common among smokers. For example, in many cases smokers are thinner than nonsmokers, tend to drink more alcohol, may be less physically active, and have poor diets. Women who smoke also tend to have an earlier menopause than nonsmokers. These factors place many smokers at an increased risk for osteoporosis apart from their tobacco use.

In addition, studies on the effects of smoking suggest that smoking increases the risk of having a fracture. As well, smoking has been shown to have a negative impact on bone healing after fracture.

Osteoporosis Management Strategies

Start by quitting. The best thing smokers can do to protect their bones is to quit smoking. Smoking cessation, even later in life, may help limit smoking-related bone loss. Many resources are available to help you stop smoking, some of which are listed at the end of this fact sheet.

Eat a well-balanced diet rich in calcium and vitamin D. Good sources of calcium include low-fat dairy products; dark green, leafy vegetables; and calcium-fortified foods and beverages. Supplements can help ensure that you get adequate amounts of calcium each day, especially in people with a proven milk allergy. The Institute of Medicine recommends a daily calcium intake of 1,000 mg (milligrams) for men and women up to age 50. Women over age 50 and men over age 70 should increase their intake to 1,200 mg daily.

Vitamin D plays an important role in calcium absorption and bone health. Food sources of vitamin D include egg yolks, saltwater fish, and liver. Many people, especially those who are older, may need vitamin D supplements to achieve the recommended intake of 600 to 800 IU (International Units) each day.

Exercise for your bone health. Like muscle, bone is living tissue that responds to exercise by becoming stronger. Weight-bearing exercise that forces you to work against gravity is the best exercise for bone. Some examples include walking, climbing stairs, weight training, and dancing. Regular exercise, such as walking, may help prevent bone loss and will provide many other health benefits.

Avoid excessive use of alcohol. Chronic alcohol use has been linked to an increase in fractures of the hip, spine, and wrist. Drinking too much alcohol interferes with the balance of calcium in the body. It also affects the production of hormones, which have a protective effect on bone, and of vitamins, which we need to absorb calcium. Excessive alcohol consumption also can lead to more falls and related fractures.

Talk to your doctor about a bone density test. A bone mineral density (BMD) test measures bone density at various sites of the body. This safe and painless test can detect osteoporosis before a fracture occurs and can predict one's chances of fracturing in the future. If you are a current or former smoker, you may want to ask your health care provider whether you are a candidate for a BMD test, which can help determine whether medication should be considered.

See if medication is an option for you. There is no cure for osteoporosis. However, several medications are available to prevent and treat the disease in postmenopausal women and in men. Your doctor can help you decide whether medication might be right for you.

Resources

For more information on osteoporosis, contact the:
**NIH Osteoporosis and Related Bone Diseases
National Resource Center**

2 AMS Circle

Bethesda, MD 20892-3676

Phone: 202-223-0344

Toll free: 800-624-BONE

TTY: 202-466-4315

Fax: 202-293-2356

Website: www.bones.nih.gov

Email: NIHBoneInfo@mail.nih.gov

If you need more information about available resources in your language or another language, please visit our website or contact the NIH Osteoporosis and Related Bone Diseases ~ National Resource Center.

BeTobaccoFree.gov

Website: <http://betobaccofree.hhs.gov>

FDA 101: Smoking Cessation Products

Website: www.fda.gov/forconsumers/consumerupdates/ucm198176.htm

Smokefree.gov

Website: www.smokefree.gov

For Your Information

This publication contains information about medications used to treat the health condition discussed here. When this publication was developed, we included the most up-to-date (accurate) information available. Occasionally, new information on medication is released.

For updates and for any questions about any medications you are taking, please contact the Food and Drug Administration toll free at 888-INFO-FDA (463-6332) or visit its website at www.fda.gov. For additional information on specific medications, visit Drugs@FDA at www.accessdata.fda.gov/scripts/cder/drugsatfda. Drugs@FDA is a searchable catalog of FDA-approved drug products.

NIH Publication No. 16-7883