

What Is Lichen Sclerosus?

Fast Facts: An Easy-to-Read Series of Publications for the Public

What Is Lichen Sclerosus?

Lichen sclerosus is a long-term problem of the skin. It mostly affects the genital and anal areas. Sometimes, lichen sclerosus appears on the upper body, breasts, and upper arms.

Who Gets Lichen Sclerosus?

Lichen sclerosus appears in:

- Women (often after menopause)
- Men (uncommon)
- Children (rare).

What Are the Symptoms?

Early in the disease, small white spots appear on the skin. The spots are usually shiny and smooth. Later, the spots grow into bigger patches. The skin on the patches becomes thin and crinkled. Then the skin tears easily, and bright red or purple bruises are common. Sometimes, the skin becomes scarred. If the disease is a mild case, there may be no symptoms.

Other symptoms are:

- Itching (very common)
- Discomfort or pain
- Bleeding
- Blisters.

What Causes Lichen Sclerosus?

Doctors don't know the exact cause of lichen sclerosus. Some doctors think a too active immune system and hormone problems may play a role. It is also thought that people inherit the likelihood of getting the disease. Sometimes, lichen sclerosus appears on skin that has been damaged or scarred from some other previous injury.

Lichen sclerosus is not contagious (it can't be caught from another person).

How Is It Diagnosed?

Doctors can look at severe lichen sclerosus and know what it is. But usually, a doctor takes a small piece of the skin patch (biopsy) and looks at it under a microscope. This allows doctors to make sure that it is not a different disease.

U.S. Department of Health
and Human Services
Public Health Service

National Institute of
Arthritis and
Musculoskeletal and
Skin Diseases
National Institutes of Health
1 AMS Circle
Bethesda, MD 20892-3675

Phone: 301-495-4484
Toll free: 877-22-NIAMS
TTY: 301-565-2966
Fax: 301-718-6366
Email:
NIAMSinfo@mail.nih.gov
Website:
www.niams.nih.gov



National Institute of
Arthritis and Musculoskeletal
and Skin Diseases

How Is It Treated?

If you have patches on the arms or upper body, they usually don't need treatment. The patches go away over time.

Lichen sclerosus of the genital skin should be treated. Even if it isn't painful or itchy, the patches can scar. This can cause problems with urination or sex. There is also a very small chance that skin cancer may develop in the patches.

Surgery is normally a good option for men. Circumcision (removing the foreskin on the penis) is the most widely used therapy for men with lichen sclerosus. The disease usually does not come back. Surgery is normally not a good option for women. When the lichen sclerosus patches are removed from the genitals of women and girls, they usually come back.

Treatment also includes using very strong cortisone cream or ointment on the skin. You put these creams on the patches every day for several weeks. This stops the itching. Then you use the cream or ointment two times a week for a long time to keep the disease from coming back. Treatment does not fix the scarring that may have already occurred.

You need regular followup by a doctor because using these creams and ointments for a long time can cause:

- Thinning and redness of the skin
- Stretch marks where the cream is applied
- Genital yeast infections.

Sometimes, you don't get better when using the cortisone creams. Some things that can keep symptoms from clearing up are:

- Low estrogen levels
- Infection
- Allergy to the medication.

When creams and ointments don't work, your doctor may suggest:

- Retinoids, or vitamin A-like drugs
- Tacrolimus ointment
- Ultraviolet light treatments (not used on skin of the genitals).

If you need medicine, ask your doctor:

- How does the medicine work?
- What are its side effects?
- Why is it the best treatment for my lichen sclerosus?

If a young girl gets lichen sclerosus, she may not require lifelong treatment. Lichen sclerosus sometimes goes away at puberty. Scarring and changes in skin color may remain.

Can People With Lichen Sclerosis Have Sex?

Women with severe lichen sclerosis in the genitals may not be able to have sex. The disease can cause scars that narrow the vagina. Also, sex can hurt and cause the patches to bleed. However, treatment with creams or ointments can help. Women with severe scarring in the vagina may need surgery, but only after lichen sclerosis is controlled with medication.

Is Lichen Sclerosis Related to Cancer?

Lichen sclerosis does not cause skin cancer. However, skin that is scarred by lichen sclerosis is more likely to develop skin cancer. If you have the disease, see the doctor every 6 to 12 months. The doctor can look at and treat any changes in the skin.

What Kind of Doctor Treats Lichen Sclerosis?

Lichen sclerosis is treated by:

- Dermatologists (doctors who treat the skin)
- Gynecologists (doctors who treat the female reproductive system)
- Urologists (doctors who treat the urinary or urogenital tract)
- Primary health care providers.

For More Information About Lichen Sclerosis and Other Related Conditions:

National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS)
Information Clearinghouse
National Institutes of Health
1 AMS Circle
Bethesda, MD 20892-3675
Phone: 301-495-4484
Toll free: 877-22-NIAMS (226-4267)
TTY: 301-565-2966
Fax: 301-718-6366
Email: NIAMSinfo@mail.nih.gov
Website: www.niams.nih.gov

For Your Information

This publication may contain information about medications used to treat the health condition discussed here. When this publication was printed, we included the most up-to-date (accurate) information available. Occasionally, new information on medication is released.

For updates and for any questions about any medications you are taking, please contact the U.S. Food and Drug Administration (FDA) toll free at 888-INFO-FDA (888-463-6332) or visit its website at www.fda.gov. For additional information on specific medications, visit Drugs@FDA at www.accessdata.fda.gov/scripts/cder/drugsatfda. Drugs@FDA is a searchable catalog of FDA-approved drug products.